# STD 20/20: A National Perspective on STD Epidemics

Jessica Frasure-Williams
Director, Programs and Partnerships
National Coalition of STD Directors



Q Search

Contact

STD INFORMATION & RESOURCES


DONATE




ncsddc.org 2

### **Outline**

- 2016 STD Surveillance Data
  - National
  - Statewide
- Strategies for STD prevention
  - Preventing infection
  - Finding and treating cases
  - Partner management strategies





#### What do you do for a living?

**Health Educator** 

Medical Provider (MD, RN, NP, MA, etc)

Field Epidemiologist

Health Department Administrator

**CBO Administrator** 


None of these. I feel

so left out.

Start the presentation to activate live content


If you see this message in presentation mode, install the add-in or get help at PollEv.com/app





# New STD cases hit record high in US, CDC says In 2016, Americans were infected with more than 2 milli... cnn.com





# The STATE of STDs in the United States



in 2016

STDS TIGHTEN THEIR GRIP ON THE NATION'S HEALTH AS RATES INCREASE FOR A THIRD YEAR



# 1.59 million CASES OF CHLAMYDIA

4.7% increase since 2015



468,514
CASES OF GONORRHEA

18.5% increase since 2015




27,814
CASES OF SYPHILIS

17.6% increase since 2015

# Annual CDC STD Prevention Budget, FY 2003– President's Proposed Budget (PB) FY 2018

50% drop in purchasing power since 2003<sup>†</sup>




# **CHLAMYDIA**



## Chlamydia Infections, 2016


Rate per 100,000 population





	# Cases	Rank
US Total	1,598,354	-
MN	22,685	39
ND	3,455	31

# Percent of Partners Treated by Strategy Among Women Diagnosed with Chlamydia in Eight Family Planning Clinic Settings in California



# **Summary of Randomized Control Trials for Expedited Partner Therapy**

Trial	Setting and Study Population	Study Design	Intervention and Control	Persistent/Recurrent Infection Rate
CDC Project 455, B2  Schillinger et al. Sex  Transm Dis 2003;30:49-56	<ul> <li>Multi-center (6 cities)</li> <li>1996- 2000</li> <li>N=1,787, 81% followed</li> <li>Women age 14-34</li> <li>Primary care, FP, teen, STD, ED</li> </ul>	Uncomplicated CT (without GC)  1.0g azithromycin DOT  21 days – 3 months  Urine PCR/LCR	PDPT to maximum 4 partners     Control = patient-referral (verbal and written)	• Control 108/726 (14.9%) • EPT 87/728 (12.0%) OR = 0.80 (0.62 - 1.05) <sup>2</sup> = 2.67, p = .102
Seattle  Golden et al. <i>NEJM</i> 2005;.352:676-685.	<ul> <li>Seattle-King Co., WA</li> <li>1998 - 2003</li> <li>N=2,751, 68% follow-up</li> <li>Male 23%, Female 77%, age ≥14 yr (mean 23 yr)</li> <li>All reporting sites: STD, FP, private, ED</li> </ul>	Uncomplicated CT     (N=2162), GC (450), or     both (139)     AZM 1.0 g + CFX 400     mg for CT, GC     Follow-up 3-19 wk     Urine NAAT (LCR or     TMA)	<ul> <li>Patient or partner pick-up of drug at 1 of 12 pharmacies</li> <li>Control = patient-referral</li> <li>DIS assistance (both arms) if patient unable/unwilling to contact partner</li> </ul>	• Control 121/931 (13.0%) • EPT 92/929 (9.9%)  RR = 0.76 (0.59 - 0.98)  2 = 4.39, p = .04  See supplemental tables for separate CT/GC outcomes
New Orleans Urethritis Trial  Kissinger et al. Clin Inf Dis 2005;41:623-9.	<ul> <li>New Orleans, LA</li> <li>2002 - 2004</li> <li>N=629, 80% behavioral FU, 30% biological FU</li> <li>Male age ≥16, median 24 yr</li> <li>STD clinic</li> </ul>	Symptomatic urethritis (61% GC, 21% CT, 6% both) AZM 1.0 g ± CFX 400mg or cipro 500 mg GenProbe(enrollment) Urine PCR (follow-up)	PDPT     Control = patient-referral with brief counseling     Third arm "Booklet Referral"	• Control 35/82 (43%) • EPT 20/87 (23%) • Booklet 16/112 (14%)  EPT v control: OR = 0.38 (0.19 – 0.74) BEPR v control: OR = 0.22 (0.11 – 0.44)

# **GONORRHEA**



## Gonorrhea Infections, 2016

Rate per 100,000 population
145.8

132.1



	# Cases	Rank
US Total	468,514	-
MN	5,104	37
ND	1,000	26




# **SYPHILIS**



### Primary & Secondary Syphilis Infections, 2016








	# Cases	Rank
US Total	24,814	-
MN	306	27
ND	2.4	33

## **Ocular Syphilis Increase**

Ocular syphilis: a manifestation of neurosyphilis affecting the eyes, which can occur at any stage of syphilis

- CDC issued a clinical advisory for ocular syphilis on 4/3/2015
- Majority of cases are among HIV-infected MSM
- 8 jurisdictions have reviewed their surveillance data, and have identified 382 cases from 2014 and 2015
- Several cases have resulted in significant outcomes including blindness


## CDC Released a Syphilis Call to Action



#### Actions included for

- Pregnant women
- MSM
- Healthcare providers
- Public Health Departments
- Decision-makers
- Community leaders
- Researchers
- EMR developers and vendors

Primary and Secondary Syphilis — Reported Cases\* by Sex, Sexual Behavior, and Race/Ethnicity, United States, 2016




# **CONGENITAL SYPHILIS**



### **Congenital Syphilis, 2016**

Rate per 100,000 Births





	# Cases	Rank
US Total	628	_
MN	7	19
ND	0	-


#### From 2015 to 2016, rates of syphilis increased:

#### **Among women**

- By 36 percent
- From 1 case per 100,000 women in 2015 to
   2 per 100,000 women in 2016

#### Among newborns (congenital syphilis)

- By 28 percent
- From 12 cases per 100,000 live births in 2015 to 16 per 100,000 live births in 2016




# Characteristics of Mothers Who Gave Birth to Infants with Congenital Syphilis in the U.S., 2014

	Number (N=458)	Percent
Did not receive prenatal care	100	21.8%
Received prenatal care (N=314, 68.6%)		
No treatment	135	29.5%
Treated <30 days prior to delivery	78	17.0%
Non-penicillin therapy	3	0.7%
Inadequate regimen for stage	13	2.8%
Adequate treatment	43	9.4%
Unknown treatment status	42	9.2%
Unknown prenatal care status	44	9.6%

# **EXTRAGENITAL TESTING**




# Most Rectal Gonorrhea Infections in MSM are Asymptomatic




# Proportion of Gonorrhea and Chlamydia Tests Missed with Urethral-only Screening among MSM

STD Surveillance Network\*, July 2010- June 2012



# Don't forget the "triple dip" for MSM





#### STD Diagnosis is an Indication of HIV Risk

Rectal GC or CT



1 in 15 MSM were diagnosed with HIV within 1 year.\*

**P&S Syphilis** 



1 in 18 MSM were diagnosed with HIV within 1 year.\*\*

# **STRATEGIES**








Visit NCSD's website to request a donation.

ncsddc.org

#### bhocpartners.org





Include a widget to search for STD testing locations right on your website!



Add this widget to your website

#### <iframe

src="https://www.cdc.gov/std/widgets/widget-6.html" name="STDwidget" id="lframe3" title="Widget for finding an STD testing site." frameborder="0" scrolling="no" height="200" width="250" marginheight="0" marginwidth="0" style="border:1px solld #ccc;">Find an STD testing site near you.

#### **IWTK I WANT THE KIT**

**ABOUT IWTK** 

WHAT IS AN STI?

TESTING

RESOURCES

RISK QUIZ

Residents of Maryland, Washington DC, and Alaska can order a kit to collect a sample in the privacy and comfort of their home, then mail it back to be tested for chlamydia, gonorrhea, and trichomonas. It's easy to do and the results are reliable!

GET THE STI KIT!



GET THE HIV KIT!



ENTER MY RESULTS



If you are interested but not eligible, registed



Home

Get a Home Test Kit

Get Test Results

Find a Clinic

The Facts

**Getting Tested** 

**Getting Treated** 

More Questions?

Call (510) 268-2110 or email Alameda@DontThinkKnow.org

> Men, click here for more information about where to get tested and treated for STDs.

Change Location



That hooking up can have a downside.

Know the Facts

About Chlamydia & Gonorrhea

Get a Home Test Kit 👂

A Free, Simple, Private Test

Get Your Test Results O

Find a Clinic

Enter zip code

FIND IT



EN ESPAÑOL







Image courtesy Complete Care Community Health Center

#### MSM Sexual Health Standards of Care

Addressing the Sexual Health Crisis among Gay, Bisexual, and Other Men who have Sex with Men (MSM)





TOPICS & ISSUES

FOR PROFESSIONALS

POLICY & ADVOCACY



LESSON PLANS

PUBLICATIONS BY TOPIC

Best Practices for Youth Friendly Clinical

TELL THEM | GET CHECKED | STD INFO | RESOURCES | TESTING



- Tell them if you've been exposed to an STD - send an anonymous ecard.
- Learn about cures and treatments.
- Find a clinic near you.

Community Guidelines -> In the News ->



Tell your hookups, ex's, boyfriends, girlfriends, and partners they may have been exposed to an STD.





SO THEY CAN KNOWBOUT US FOR PROVIDERS

# Let your partners know they should get tested

Send an anonymous email, contact your health department, or get tips on talking to your partners

GET STARTED +







PROTECT and your partners.






TEST



## **Summary**


- STDs are all increasing!
- Pay attention to emerging issues
  - Congenital Syphilis
  - Gonorrhea Resistance
  - Extragenital Testing for MSM and PrEP
- Strategies
  - Health Promotion and Condom Access
  - Linkages to Testing and Competent Care
  - Partner Notification

### **Become a Member of NCSD**







# Join us for the NCSD Annual Meeting



ncsddc.org 42







#### Jessica Frasure-Williams

Director, Programs and Partnerships

jfrasure-williams@ncsddc.org

office: 202-715-3861 mobile: 510-579-1657

